



THE UNIVERSITY OF
MELBOURNE

**Age-standardised mortality and cause of death in
the Latrobe Valley at the time of (and five years prior to)
the Hazelwood coalmine fire in Morwell, Victoria**

**Louisa Flander PhD
Driss Ait Ouakrim PhD, MPH
Seyedeh Ghazaleh Dashti DDS, MPH
Antony Ugoni MSc**

**Centre for Epidemiology & Biostatistics
Melbourne School of Population & Global Health**

l.flander@unimelb.edu.au

30 May 2015

Executive Summary

This report examines mortality data from the Latrobe Valley postcodes during the period of the Hazelwood coalmine fire in February and March 2014. The mine fire occurred during a period of high temperatures with associated health risks (<http://www.dpc.vic.gov.au/index.php/news-publications/hazelwood-mine-fire-inquiry-report>, Hazelwood Mine Fire Report 2014). The mortality data are compared to the same summer months in the previous five years to examine this association. The mortality from the period February to June 2014 is included also in this report, and compared to the same period in the previous five years. These comparisons were made to examine whether there were associations of the Hazelwood fire on mortality beyond the summer months when the fire occurred.

The analyses examine these associations in terms of the statistical evidence linking the deaths with the occurrence of the mine fire. The findings reported are based on the small number of deaths in the affected postcodes, which limits the interpretation of the results.¹ Key findings in this report are to be interpreted cautiously, with the understanding that moderate or no statistical evidence of association cannot be interpreted as evidence for or against a particular cause of death.

The analysis of these data shows no statistical evidence that 2014 mortality rates differ from comparable rates for the same months in 2009, a season similar to 2014 with high temperatures and particulate matter from bushfire smoke. Broad confidence intervals for the 2009–13 rate ratios, which approach or overlap the confidence intervals of the 2014 rates, express the lack of statistical evidence for an overall difference in the mortality rates throughout the study period.

There is statistical evidence that air quality exceedances are associated with mortality throughout the study period, not just during the period of the 2014 Hazelwood coalmine fire, or the 2009 bushfire.² Overall for 2009–14, most deaths associated with air quality exceedances in the affected postcodes occurred outside of the February–March period; 85% of these occurred in 2012 and 2013.

Mortality in all age groups was 2.13 times higher on days with air quality exceedances, compared to days without air quality exceedances for the period 2009–

¹ The associations reported herein are given as point estimates of rates with associated 95% confidence intervals. The 95% confidence intervals given with a point estimate is equivalent to the statement that there is a 95% probability that the value of the point estimate lies within the stated range of values. These intervals can be broader or narrower depending on several factors, including sample size and population variability. When the confidence interval contains one (1), the evidence for an association is weak. We note that non-significant results in the case of small sample sizes such as those in this report are prone to misinterpretation, leading to the conclusion of an effect where there is none, or the conclusion of no effect where there is one (see Altman DG and Bland JM, 1995, Absence of evidence is not evidence of absence, *British Med J* 311:485).

² An exceedance is an instance or condition where the observed concentration of a pollutant goes beyond the permitted quality standard or threshold. The threshold level for this analysis is the daily mean value of 50µg/m³, although particulate matter is a non-threshold pollutant (<http://www.epa.vic.gov.au/your-environment/air/bushfires-and-air-quality>, see also Table 1, Standards and goals for pollutants other than particles as PM_{2.5}, National Environment Protection (Ambient Air Quality) Measure, <http://www.comlaw.gov.au/Details/C2004H03935>).

14 ($p < 0.01$). The mortality in the vulnerable age group 65 years and older was 2.0 times higher on exceedance days compared to days without exceedances for the period ($p < 0.01$).

There is no statistical evidence for the association of daily average temperature at or over 30° C with mortality in the February-March period for 2009–14. There is moderate evidence that colder temperatures are associated with mortality in the February-June period for 2009–14.

draft

Methods

Births, Deaths, and Marriages Victoria provided the mortality data for these analyses in the form of Stata files containing all Victorian deaths for the period 2009–2014. For each death, the dataset included variables for date of death, age, 5-year age group, sex, cause of death and postcode.

There were 3 414 deaths in the Latrobe Valley postcodes for the years 2009–2014. Our analysis is based on the 3 398 ascertained deaths. There were 13 deaths listed as ‘unascertained’; of these, there were two unascertained deaths in 2014 (May and September). Three additional cases were eliminated from the final analysis due to missing data in other variables.

Cause of death categories and definitions

Cause of death for the total Victorian 203 963 deaths for 2009–14 was given in the form of text description of the underlying cause of death. Using the *regular expressions* command in Stata 13.0, we generated variables that identified deaths known to be associated with exposure to airborne particulate matter and/or pollutants. These categories are deaths by *respiratory* conditions, *cardiovascular* conditions, and deaths with *direct relationship to fire*.

Causes of death due to *respiratory conditions* included chronic obstructive pulmonary disease, asthma, pneumonia, bronchitis, bronchopneumonia, pulmonary embolism, pulmonary fibrosis, pulmonary oedema, and respiratory arrest.

Causes of death due to *cardiovascular conditions* included myocardial infarction, ischemic heart disease, congestive heart failure, coronary heart disease, cardiomyopathy, aortic dissection, aortic stenosis, arterial fibrillation, ventricular fibrillation, cardiac amyloidosis, cardiac arrhythmia and tachycardia, and cardiac arrest.

Due to the small number of deaths in the four postcodes of interest, the aggregated variable *cardiorespiratory* conditions for causes of death due to respiratory *and/or* cardiovascular conditions was generated by combining the above two variables.

Causes of death from *direct relationship with fire* included carbon monoxide poisoning, inhalation of smoke and fire gases, complications of thermal burn injuries, and general effects of fire.

Other causes of death included accidents or injuries, suicide, drug and alcohol toxicity, sepsis or infection, cancer, failure or disease of systems other than the respiratory and cardiovascular systems. In these analyses the number of deaths due to *respiratory*, *cardiovascular* or *cardiorespiratory* causes was compared with number of deaths due to *other causes of death*.

Temperature and air quality variables

The Morwell Bureau of Meteorology Site 85280 at the Latrobe Valley Airport, closely located to the four Latrobe Valley postcodes in this analysis, provided daily mean temperatures for 2009–14. The threshold level for this analysis is daily mean temperature in excess of 30° Celsius, which triggers the state’s Heat Health Alert System.

The Environmental Protection Agency Traralgon air quality monitoring site, closely located to the four Latrobe Valley postcodes in this analysis, provided daily mean measures of particulate matter in excess of 10 micrometers or less in diameter (PM₁₀) for 2009-14. The threshold level used in our analysis is 50µg/m³, although particulate matter is a non-threshold pollutant (<http://www.epa.vic.gov.au/your-environment/air/bushfires-and-air-quality>, see also Table 1, Standards and goals for pollutants other than particles as PM_{2.5}, National Environment Protection (Ambient Air Quality) Measure, <http://www.comlaw.gov.au/Details/C2004H03935>).

Age-standardisation

Age-standardisation allows for comparison of mortality rates over the time period 2009–2014. This is done by adjusting each year's deaths to account for any possible age differences in the structure of the local population. For example, in the case of the four Latrobe Valley postcodes, there is a five-fold difference in the population size of Churchill compared to Traralgon, and a nine-year difference in the median age of Churchill compared to Morwell (<http://www.abs.gov.au/websitedbs/censushome.nsf/home/Census?opendocument#from-banner=GT>).

Due to the small population size of the Latrobe Valley, the Morwell, Churchill, Moe and Traralgon postcodes' mortality data were aggregated for age standardisation into three age-categories, under age 50 years, 50-64 years and 65 years and over. For age-standardisation we used the direct method (Australian Institute of Health and Welfare 2011. Principles on the use of direct age-standardisation in administrative data collections: for measuring the gap between Indigenous and non-Indigenous Australians. Cat. No. CSI 12. Canberra: AIHW). We created age-specific population estimates from the Australian Bureau of Statistics using the age distribution of the 2011 Australian standard population (<http://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/3101.0Sep%202014?OpenDocument>).

Multivariable modelling

Poisson regression models adjusted for age and sex distribution and for temperature and air quality in excess of threshold values were constructed to calculate mortality rate ratios and associated 95% confidence intervals. These rate ratios were used to compare the mortality observed for the periods February-June and February-March 2014 with the mortality observed during those periods for each year from 2009 to 2013. The 2014 mortality rates were used as the reference rates (mortality rate ratio of 1.0).

These analyses were conducted for *all-cause* mortality and mortality due to *respiratory*, *cardiovascular* and the combined category of *cardiorespiratory* causes. Mortality rate ratios for the aggregated vulnerable age groups aged 65 years and older were examined for all-cause mortality, and *respiratory*, *cardiovascular* and *cardiorespiratory* causes. There were insufficient cases to analyse the vulnerable age groups aged 5 years and younger.

Results

Age-adjusted mortality

The difference in deaths between 2014 and the previous five years can be seen in the age-standardised mortality rates (Table 1, Table 2). These tables show that the mortality rates for these postcodes were similarly elevated for 2009 and 2014, compared to the years 2010–13, when rates were almost uniformly lower. This is best seen in the comparison between February-March, 2009 and 2014, when the age-standardised rates are 1.5 deaths per 1000 person-years³ and 1.6 deaths per 1000 person-years respectively (Table 2). Deaths in February-March, 2010–2013 were 1.1 to 1.2 per 1000 person-years. These results should be interpreted with caution, as the comparisons are not statistical evidence of differences between each year's mortality.

Table 1. Age-standardised* mortality rates (ASR) in the Latrobe Valley per 1,000 person-years between February-June, 2009–2014**

Age category	2009		2010		2011		2012		2013		2014	
	<i>n</i>	<i>ASR</i>	<i>n</i>	<i>ASR</i>	<i>n</i>	<i>ASR</i>	<i>n</i>	<i>ASR</i>	<i>n</i>	<i>ASR</i>	<i>n</i>	<i>ASR</i>
< 50	24	0.3	18	0.4	12	0.3	18	0.2	29	0.3	18	0.4
50-64	22	0.4	24	0.4	34	0.4	32	0.5	27	0.3	32	0.3
≥ 65	225	2.4	189	2.1	184	2.0	157	2.3	170	2.4	188	2.8
All ages	271	3.4	231	3.2	230	3.0	207	3.3	226	3.3	238	3.9

*Directly age-standardised using the 2011 Australian standard population

**Latrobe Valley defined as Morwell (3840), Churchill (3842), Moe (3825) and Traralgon (3844).

Table 2. Age-standardised* mortality rates (ASR) in the Latrobe Valley per 1,000 person-years between February-March, 2009–2014**

Age category	2009		2010		2011		2012		2013		2014	
	<i>n</i>	<i>ASR</i>	<i>n</i>	<i>ASR</i>	<i>n</i>	<i>ASR</i>	<i>n</i>	<i>ASR</i>	<i>n</i>	<i>ASR</i>	<i>n</i>	<i>ASR</i>
< 50	7	0.1	6	0.2	4	0.1	8	0.1	13	0.1	8	0.1
50-64	10	0.2	9	0.1	10	0.2	14	0.1	9	0.1	16	0.1
≥ 65	94	1.1	63	0.7	68	0.7	59	0.9	59	0.8	84	1.2
All ages	111	1.5	78	1.2	82	1.2	81	1.2	81	1.1	108	1.6

*Directly age-standardised using the 2011 Australian standard population

**Latrobe Valley defined as Morwell (3840), Churchill (3842), Moe (3825) and Traralgon (3844).

³ The person-year is a measure of the estimated time-at-risk for the population under review.

Temperature and air quality

Comparison of the mortality records with the environmental observations shows that there were more deaths associated with extreme temperatures in 2009 and 2014 than in the years 2010–13 (Table 3). There were 27 deaths that occurred on days with mean temperature at or over 30° C in the affected postcodes for these two years, 13 in 2009 and seven that occurred in 2014. About half of the 93 deaths that occurred on days with mean air quality at or over $\geq 50\mu\text{g}/\text{m}^3$ PM₁₀ in the affected postcodes for 2009–14 occurred in 2013; nine occurred in 2014.

Comparison of these records for February-March, 2009–2014, shows that deaths on days with air quality levels $\geq 50\mu\text{g}/\text{m}^3$ PM₁₀ were similarly high in 2009 and 2014 (Table 4). For 2009 and 2014, 67% of the deaths on days with average PM₁₀ levels $\geq 50\mu\text{g}/\text{m}^3$ occurred during the fire months of February-March, compared to deaths occurring with similar exposures at other times of the year. There were three deaths in 2011, and all were associated with air quality levels $\geq 50\mu\text{g}/\text{m}^3$ PM₁₀ in the February-March period (100%).

We note that 68% of the total 93 deaths for February-June 2009–14 deaths occurred on days of air quality exceedances in 2012 and 2013, with most occurring outside the months of February-March (compare Tables 3, 4). Overall for 2009–14, most deaths associated with air quality exceedances in the affected postcodes occurred outside of the February-March period during the months April-June; 85% of these occurred in 2012 and 2013, during the months April-June.

Table 3. Latrobe Valley* annual number of deaths associated with temperature or air quality exceedances, February-June, 2009–2014

	2009	2010	2011	2012	2013	2014	Total
Temperature $\geq 30^\circ\text{C}$	13	4	0	0	3	7	27
Air quality $\geq 50\mu\text{g}/\text{m}^3$ PM₁₀	15	3	3	17	46	9	93

*Latrobe Valley defined as Morwell (3840), Churchill (3842), Moe (3825) and Traralgon (3844).

Table 4. Latrobe Valley* number of deaths associated with air quality exceedances, February- March, 2009–2014

	2009	2010	2011	2012	2013	2014	Total
Air quality $\geq 50\mu\text{g}/\text{m}^3$ PM₁₀	10	2	3	0	8	6	27

*Latrobe Valley defined as Morwell (3840), Churchill (3842), Moe (3825) and Traralgon (3844).

Cause of death (Tables 5-8, Figure 1)

Analyses of cause of death are to be interpreted cautiously, with the understanding that moderate or no statistical evidence of association cannot be interpreted as evidence for or against a particular cause of death.

There were 10 deaths from *direct relationship with fire* between 2009 and 2014 in the Latrobe Valley; of these, six deaths occurred in February-June 2009 and one in February-June 2013. No deaths from *direct relationship with fire* occurred in February-June 2010–12 or 2014.

We found moderate statistical evidence for a 16% lower *all-cause* mortality rate for February-June 2013 compared to the same period in 2014 (Table 5, $p=0.02$). There was no statistical evidence for any other differences in mortality for that period in 2009–12 compared to 2014.

During the period February-March, the *all-cause* mortality rate was 31% lower in 2013 ($p=0.01$) and 24% lower in 2012 ($p=0.05$) compared to the same period in 2014 (Table 5). There was no statistical evidence for other differences in mortality for this period in 2009–11. The ratio of the mortality rates of previous years to the mortality rates of 2014 has broad and overlapping associated 95% confidence intervals (Table 5, Figure 1).

Air quality exceedances for the entire period were associated with *all-cause* mortality throughout this period (Table 5). Mortality in the February-March period was 2.13 times higher on days with air quality $\geq 50\mu\text{g}/\text{m}^3$ PM_{10} compared to less extreme days ($p<0.01$). Mortality in the February-June period was 1.83 times higher ($p<0.01$).

Temperature exceedances do not show statistical evidence of association with *all-cause* mortality in the February-March period 2009–13 compared to February-March 2014. We note that there is moderate statistical evidence for the association of colder temperatures with February-June *all-cause* mortality for all ages.

Cardiovascular mortality for all ages was 42% lower in 2009 compared to 2014, for the February-March period, after adjusting for age, sex, mean daily temperature and 24-hour air quality (Table 7, $p=0.05$). There was no statistical evidence for differences in mortality rates due to the other smoke exposure causes (*respiratory* and combined *cardiorespiratory* causes) for these time periods (Table 6, Table 8).

Table 5. Latrobe Valley* *all-cause* mortality in 2009–2013 compared to 2014

Year	February-June			February-March		
	Rate Ratio	(95% CI)	p-value	Rate Ratio	(95% CI)	p-value
Year						
2014	1	–	–	1	–	–
2013	0.84	(0.74-0.97)	0.02	0.69	(0.52-0.90)	0.01
2012	0.87	(0.76-1.00)	0.06	0.76	(0.59-1.00)	0.05
2011	0.89	(0.78-1.03)	0.13	0.86	(0.67-1.12)	0.29
2010	0.95	(0.83-1.09)	0.52	0.84	(0.65-1.09)	0.20
2009	0.93	(0.81-1.06)	0.30	1.01	(0.79-1.28)	0.91
Temperature						
< 30° C	1	–	–	1	–	–
≥ 30° C	0.55	(0.34-0.89)	0.02	1.24	(0.78-1.97)	0.35
PM₁₀						
< 50ug/m ³	1	–	–	1	–	–
≥ 50ug/m ³	1.83	(1.57-2.14)	<0.01	2.13	(1.55-2.91)	<0.01

*Latrobe Valley defined as Morwell (3840), Churchill (3842), Moe (3825) and Traralgon (3844). P-value is the probability of observing a rate ratio this small or smaller given there was no difference in rates compared to 2014.


Figure 1. All cause mortality rate ratios in the Latrobe Valley, 2009–2013 compared to 2014 (Reference rate 1.0)

Table 6. Mortality due to respiratory causes in the Latrobe Valley* in 2009–13 compared to 2014

Year	February-June			February -March		
	Rate Ratio	(95% CI)	p-value	Rate Ratio	(95% CI)	p-value
2014	1	-	-	1	-	-
2013	1.17	(0.78-1.76)	0.43	1.02	(0.49-2.15)	0.94
2012	1.04	(0.68-1.59)	0.84	1.61	(0.83-3.12)	0.16
2011	1.43	(0.96-2.13)	0.08	1.59	(0.82-3.07)	0.16
2010	1.43	(0.97-2.12)	0.07	1.27	(0.63-2.57)	0.49
2009	0.95	(0.61-1.47)	0.82	1.08	(0.54-2.17)	0.81

*Latrobe Valley defined as Morwell (3840), Churchill (3842), Moe (3825) and Traralgon (3844). P-value is the probability of observing a rate ratio this small or smaller given there was no difference in rates compared to 2014.

Table 7. Mortality due to cardiovascular causes in the Latrobe Valley* in 2009–13 compared to 2014

Year	February-June			February-March		
	Rate Ratio	(95% CI)	p-value	Rate Ratio	(95% CI)	p-value
2014	1	-	-	1	-	-
2013	0.80	(0.57-1.12)	0.21	0.77	(0.46-1.28)	0.32
2012	0.86	(0.62-1.20)	0.39	0.59	(0.34-1.05)	0.08
2011	0.79	(0.56-1.14)	0.22	0.65	(0.38-1.13)	0.14
2010	0.84	(0.60-1.19)	0.34	0.60	(0.34-1.07)	0.09
2009	0.70	(0.49-1.00)	0.06	0.58	(0.34-0.99)	0.05

*Latrobe Valley defined as Morwell (3840), Churchill (3842), Moe (3825) and Traralgon (3844). P-value is the probability of observing a rate ratio this small or smaller given there was no difference in rates compared to 2014.

Table 8. Mortality due to cardiorespiratory causes in the Latrobe Valley* in 2009–13 compared to 2014

Year	February-June			February-March		
	Rate Ratio	(95% CI)	p-value	Rate Ratio	(95% CI)	p-value
2014	1	-	-	1	-	-
2013	0.95	(0.74-1.22)	0.72	0.89	(0.60-1.33)	0.6
2012	0.97	(0.76-1.24)	0.83	0.94	(0.63-1.39)	0.77
2011	1.07	(0.84-1.37)	0.55	0.99	(0.67-1.45)	0.97
2010	1.06	(0.83-1.35)	0.63	0.84	(0.56-1.28)	0.44
2009	0.81	(0.62-1.05)	0.14	0.78	(0.52-1.15)	0.22

*Latrobe Valley defined as Morwell (3840), Churchill (3842), Moe (3825) and Traralgon (3844). P-value is the probability of observing a rate ratio this small or smaller given there was no difference in rates compared to 2014.

Mortality in the vulnerable age group 65 years and older (Tables 9-12, Figure 2)

Analyses of cause of death in this age group are to be interpreted cautiously, with the understanding that moderate or no statistical evidence of association cannot be interpreted as evidence for or against a particular cause of death.

We found moderate statistical evidence for a 15% lower *all-cause* mortality rate for February-June 2012 compared to the same period in 2014 (Table 9, $p=0.04$) for the vulnerable age group 65 years and older, after adjusting for age, sex, mean daily temperature and 24-hour air quality. The mortality rate for the February-March 2013 period was 32% lower for this age group compared to the same period in 2014 (Table 9, $p=0.01$). The ratio of the mortality rates for 2009–13 to the mortality rates of 2014 has broad and overlapping associated 95% confidence intervals and must be interpreted with caution (Table 9, Figure 2).

Air quality exceedances for the entire period were associated with *all-cause* mortality throughout this period for this age group (Table 9). Mortality in the February-March period was 2.0 times higher on days with air quality $\geq 50\mu\text{g}/\text{m}^3$ PM_{10} compared to less extreme days ($p<0.01$). Mortality in the February-June period was 1.74 times higher ($p<0.01$).

Temperature exceedances do not show statistical evidence of association with *all-cause* mortality in the February-March period 2009–13 compared to February-March 2014 for this age group. We note that there is moderate statistical evidence for the association of colder temperatures with February-June mortality for this age group.

Respiratory mortality for this age group was 57% higher in February-June 2011 compared to the same period in 2014, after adjusting for age, sex, mean daily temperature and 24-hour air quality (Table 10, $p=0.03$).

Cardiovascular mortality for this age group was 36% lower in 2009 compared to 2014, for the February-March period, after adjusting for age, sex, mean daily temperature and 24-hour air quality (Table 11, $p=0.03$). There was no statistical


evidence for differences in mortality rates due to combined *cardiorespiratory* causes for these time periods for this age group (Table 12).

Table 9. Latrobe Valley[†] all cause mortality in 2009–2013 compared to 2014, people age 65 years and older

	February-June			February-March		
	Rate Ratio	(95% CI)	p-value	Rate Ratio	(95% CI)	p-value
Year						
2014	1	–	–	1	–	–
2013	0.86	(0.74-1.00)	0.06	0.68	(0.50-0.92)	0.01
2012	0.85	(0.73-0.98)	0.04	0.75	(0.56-1.00)	0.06
2011	0.88	(0.75-1.02)	0.11	0.79	(0.59-1.07)	0.14
2010	0.93	(0.80-1.09)	0.41	0.78	(0.58-1.05)	0.11
2009	0.95	(0.82-1.10)	0.5	1.01	(0.77-1.32)	0.92
Temperature						
< 30°C	1	–	–	1	–	–
≥ 30°C	0.23	(0.07-0.79)	0.02	0.55	(0.17-1.77)	0.32
PM₁₀						
< 50ug/m ³	1	–	–	1	–	–
≥ 50ug/m ³	1.74	(0.46-2.09)	<0.01	2.00	(1.36-2.95)	<0.01

*Latrobe Valley defined as Morwell (3840), Churchill (3842), Moe (3825) and Traralgon (3844). P-value is the probability of observing a rate ratio this small or smaller given there was no difference in rates compared to 2014.

Figure 2. All cause mortality rate ratios in the Latrobe Valley, 2009–2013 compared to 2014, people age 65 years and older, (Reference rate 1.0)



DRAFT

Table 10. Mortality due to respiratory causes in the Latrobe Valley[†] in 2009–2013 compared to 2014, *people age 65 years and older*

Year	February-June			February-March		
	Rate Ratio	(95% CI)	p-value	Rate Ratio	(95% CI)	p-value
2014	1	-	-	1	-	-
2013	1.31	(0.87-1.98)	0.19	1.10	(0.53-2.28)	0.79
2012	1.10	(0.71-1.70)	0.67	1.50	(0.76-2.95)	0.23
2011	1.57	(1.03-2.38)	0.03	1.85	(0.94-3.63)	0.07
2010	1.38	(0.90-2.10)	0.13	1.02	(0.47-2.21)	0.95
2009	0.89	(0.56-1.43)	0.65	0.88	(0.41-1.84)	0.74

*Latrobe Valley defined as Morwell (3840), Churchill (3842), Moe (3825) and Traralgon (3844). P-value is the probability of observing a rate ratio this small or smaller given there was no difference in rates compared to 2014.

Table 11. Mortality due to cardiovascular causes in the Latrobe Valley[†] in 2009–2013 compared to 2014, *people age 65 years and older*

Year	February-June			February-March		
	Rate Ratio	(95% CI)	p-value	Rate Ratio	(95% CI)	p-value
2014	1	-	-	1	-	-
2013	0.87	(0.61-1.24)	0.45	0.89	(0.53-1.50)	0.68
2012	0.84	(0.59-1.20)	0.36	0.64	(0.36-1.15)	0.14
2011	0.82	(0.56-1.21)	0.33	0.64	(0.34-1.19)	0.17
2010	0.83	(0.57-1.21)	0.36	0.74	(0.41-1.32)	0.31
2009	0.64	(0.43-0.96)	0.03	0.60	(0.34-1.05)	0.08

*Latrobe Valley defined as Morwell (3840), Churchill (3842), Moe (3825) and Traralgon (3844). P-value is the probability of observing a rate ratio this small or smaller given there was no difference in rates compared to 2014.

Table 12. Mortality due to cardiorespiratory causes in the Latrobe Valley[†] in 2009–2013 compared to 2014, people age 65 years and older

Year	February-June			February-March		
	Rate Ratio	(95% CI)	p-value	Rate Ratio	(95% CI)	p-value
2014	1	-	-	1	-	-
2013	1.05	(0.82-1.35)	0.65	1.00	(0.68-1.48)	0.98
2012	0.99	(0.76-1.28)	0.94	0.96	(0.65-1.42)	0.85
2011	1.14	(0.88-1.47)	0.30	1.05	(0.70-1.57)	0.79
2010	1.04	(0.80-1.35)	0.73	0.84	(0.54-1.31)	0.46
2009	0.76	(0.57-1.02)	0.07	0.71	(0.47-1.09)	0.12

*Latrobe Valley defined as Morwell (3840), Churchill (3842), Moe (3825) and Traralgon (3844). P-value is the probability of observing a rate ratio this small or smaller given there was no difference in rates compared to 2014.

Discussion

The interpretation of the Latrobe Valley mortality data for the period of the Hazelwood coalmine fire depends on comparing these deaths with cases in the same place and at the same season in previous years. There are important caveats to note when considering the small number of deaths for these comparisons.

The results demonstrate the uncertainties around estimates for 2014 mortality in the Latrobe Valley postcodes, as the numbers are small even with aggregating the four postcodes. Estimated rates for each year and cause of death category are associated with broad and overlapping confidence intervals. Statistical evidence for or against associations with exposures to environmental factors must be interpreted against the fact of small numbers of deaths.

All-cause and specific causes of death were considered separately, as *cardiovascular* and/or *respiratory* mortality are better indicators of the effects of exposure to smoke and particulate matter. However, in these data, there were insufficient numbers of such deaths to conduct any meaningful comparison between the periods of interest in 2009–13 and 2014. Findings within the specific cause of death categories are to be interpreted with caution.

The same caveat exists for demonstrating the association of exposure to particulate matter from smoke on *all-cause* mortality in the Latrobe Valley. Whilst there were six deaths in the affected postcodes on days with air quality exceedances during the 2014 coalmine fire, there were ten such deaths in 2009 during the same period and eight such deaths in 2013 during the same period. Overall for 2009–14, most deaths associated with air quality exceedances in the affected postcodes occurred outside of the February-March period; 85% of these occurred in 2012 and 2013.

These observations mean that there is statistical evidence that air quality exceedances are associated with mortality throughout the study period, not just during the period of the Hazelwood coalmine fire. The small number of deaths restricts the analysis to air

quality measures on the date of death; it is not possible to analyse each death in association with air quality on the day, week or month before that death.

We note in this regard that air quality records for monitoring stations in the affected postcodes show that the mean daily PM_{2.5} threshold was exceeded during the February-March 2014 period except in Moe. Whilst we cannot compare these records with the same period in previous years, it does suggest that smoke exposure was variable throughout the Latrobe Valley and there may be associated differences in regional mortality that cannot be captured in our analysis.

Whilst extreme summer temperatures have been associated with increased mortality, we have no statistical evidence for this association with mortality in this dataset, once we have adjusted for the effects of air quality. The January 2014 Victorian heatwave may have affected vulnerable people in the Latrobe Valley who later died during the period of the coalmine fire. However, the small number of deaths in the affected postcodes restricts the analysis to temperatures on the date of death; it is not possible to analyse each death in association with temperatures on the day, week or month before that death.

We note that there is moderate statistical evidence for the association of colder temperatures with February-June mortality for all ages, and for the vulnerable age group 65 years and older. This may explain the 57% excess mortality due to *respiratory* causes in 2011 compared to 2014 in the vulnerable elderly. Statistical evidence of the association of colder temperatures and air quality $\geq 50\mu\text{g}/\text{m}^3$ PM₁₀ with mortality could not be demonstrated with these data; however, this lack of evidence does not rule out the possibility of such an effect.

There is moderate statistical evidence that *cardiovascular* mortality was higher during the period of the 2014 fire compared to the 2009 fire. This finding must be interpreted with caution due to the small number of deaths in these categories. There are not sufficient data to associate these excess deaths with specific extremes in air quality or temperature. However, the proposed prospective study that will track Latrobe Valley residents who were exposed during the Hazelwood fire may contribute useful information about the association of exposure to brown coal particulate matter with cardiovascular health.