

" RAL-4 "



THE UNIVERSITY OF
MELBOURNE

**Review of Birth Deaths & Marriages Victoria (BDMV)
mortality data for the Latrobe Valley and the time of
the Hazelwood coal mine fire in Morwell**

**Louisa Flander PhD
Senior Research Fellow**

**Dallas English PhD
Professor**

**Centre for Epidemiology & Biostatistics,
Melbourne School of Population & Global Health
University of Melbourne, VIC 3010**

l.flander@unimelb.edu.au

Executive Summary

Our review of the BDMV mortality data (2009-14) for the Latrobe Valley shows that slightly more deaths occurred in the period January to June 2014 compared with the period January to June 2009-13 but the evidence that this is not due to just chance alone is inconclusive.¹

This assessment reviews the data provided, and shows several ways of demonstrating the small increase in Latrobe Valley deaths in 2014, compared with 2009-13. We have shown these results using the following analyses, looking at deaths per month/all postcodes, and annual deaths/all postcodes. We have analysed reported mortality from Morwell as well.

We do not find this increase to be conclusive evidence of any particular effect, given the very wide confidence intervals around the observations, and the lack of useful denominators to compare health events in these postcodes. These uncertainties include, but may not be limited to, the small population size under review, and the fact that we have no information about the underlying age or sex distribution or population movements over time within the postcodes concerned.²

Monthly mortality

The graphic representation of reported deaths by month and associated exact confidence intervals for these observations over the period 2009-14 shows that 2014 deaths are within the range observed for the previous five years, with postcode data combined for analysis (Figure 1).³

Poisson regression

The 2009-13 data were modelled using Poisson regression (for categorical data) and the 2014 deaths were predicted using this model (see Figure 2 and Table 1). This model shows that there are 37 additional deaths overall in the 2014 period (339 observed for 2014, compared to the 302 annual average predicted by the model). The additional deaths observed occurred in March and May of 2014.

Linear Regression

The monthly number of deaths was approximately normally distributed, so the data were analysed also using linear regression. Separate analyses were done for all Latrobe Valley postcodes and for Morwell (3840). There was no evidence of

¹ Excess mortality means the number of deaths observed exceed the number predicted by statistical estimation.

² As we lack information on the population size and thus person/years (estimated time-at-risk for the population) for each postcode, we are unable to adjust for the difference in postcode population size and any subsequent difference in postcode death rate. By aggregating the postcodes into a single group (and assuming the overall population stayed relatively stable over the study period), we assume then that the number of deaths represent a single population for analysis.

³ Exact confidence interval from D. Clayton and M. Hills, *Statistical Models in Epidemiology*, Oxford University Press, Oxford, 1993. The confidence bounds are derived from the observations in this (Poisson) distribution, without assuming and thus approximating the Normal distribution.

serial autocorrelation of the residuals. 'Exposure' for this analysis was defined in two ways:

1. January-June 2014 versus January-June all other years;
2. February-March 2014 versus February-March all previous years.

For Morwell, there was weak evidence of additional deaths in 2014 compared with other years (Table 2), but for February-March, there were fewer deaths than in previous years. Wide confidence intervals (including zero) around these estimates and large p-values indicate that the results are inconclusive, since the results are consistent with large excesses in deaths in 2014 and large decreases for the February-March period. For the whole Latrobe Valley, there were 7.4 additional deaths per month in 2014 when compared with 2009-13 and 9.2 additional monthly deaths for February-March 2014 compared with February-March 2009-13. Again, the confidence intervals were wide and included zero (see Table 2), and the results inconclusive.

Limitations

We cannot conclude that the 2014 mortality observed is due to any single cause, or whether it has occurred by chance alone. We did not take external factors such as local weather conditions into account in these analyses. Analysis of the cause of deaths for this period would be required to explore common risk factors. We have no information on the underlying age/sex distribution of these localities, or of the recent demographic changes in these communities, both trends that could underlie the mortality observed in 2014.

Figure 1. Monthly mortality and associated exact 95% confidence intervals for aggregated Latrobe Valley postcodes, 2009-14




Figure 2. Monthly mortality and associated exact 95% confidence intervals, Latrobe Valley postcodes 2009-13, and Poisson predicted values, 2014




Table 1. Predicted additional monthly deaths during 2014 from Poisson regression analysis, for aggregated postcodes

	Observed	Predicted	Lower bound	Upper bound
Jan 2014	54	51.94	46.53	58.56
Feb 2014	50	43.38	38.47	48.91
Mar 2014	62	52.98	47.47	59.15
Apr 2014	54	49.90	44.70	55.70
May 2014	62	52.98	47.47	59.15
Jun 2014	57	50.40	45.15	56.26
TOTAL	339	301.58	269.78	337.72

Table 2. Predicted additional monthly deaths during 2014, compared with 2009-13, for aggregated Latrobe Valley postcodes, and for Morwell, linear regression analysis

Time period	Predicted additional deaths per month,		p-value ⁴	R ²
	Number	95% confidence interval		
All postcodes				
Jan-June 2014 vs Jan-June 2009-13	7.4	-0.69, 15.55	0.07	0.09
Feb-Mar 2014 vs Feb-Mar 2009-13	9.2	-5.48, 23.88	0.19	0.16
Morwell				
Jan-June 2014 vs Jan-June 2009-13	1.4	-2.8, 5.7	0.50	0.01
Feb-Mar 2014 vs Feb-Mar 2009-13	-2.6	-11.5, 06.3	0.53	0.04

⁴ These p-values can be interpreted as providing only weak evidence against the null hypothesis of no difference in number of deaths in 2014 versus the previous years. See B.R. Kirkwood and J.A.C. Sterne, Essential Medical Statistics, 2nd edition, Blackwell Science, Oxford, 2003.